



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

process is described by MOLISCH in a little pamphlet, entitled *Das Warmbad*.⁷ The treatment consists merely in immersing the shoots of potted plants (by inversion) in water of 30–35° C. for 9–12 hours and then keeping them in a warm (25°), moist, dark chamber until the leaves begin to appear, when further development proceeds under ordinary greenhouse conditions. Lilacs and spiraeas treated to the warm bath about mid-November flowered before Christmas, and azaleas early in January; while untreated plants, under the same conditions, had at this time hardly started their buds. The exact duration and temperature of the bath for securing the best results vary with different species and races. The process is already in use and is likely to be extensively practiced.—C. R. B.

The flora of Korea.—The first part of this work⁸ includes the families Ranunculaceae to Dipsaceae, and their sequence is essentially that of BENTHAM and HOOKER in the *Genera plantarum*. Concise dichotomous keys to the genera introduce each family containing more than one genus, and similar keys at the beginning of each genus precede the enumeration of species. Under the species a very full bibliography is given, as well as the citation of exsiccatae and the general distribution. The Japanese name is also given in many cases. Several new species and varieties are published, and the text is augmented by fifteen full-page clean-cut illustrations. An index to genera mentioned in the flora concludes the part.—J. M. GREENMAN.

North American Flora.—Volume XVII, Part I, of this work⁹ contains the following groups: Typhaceae by P. WILSON; Sparganiaceae, Elodeaceae, and Hydrocharitaceae by P. A. RYDBERG; Zannichelliaceae, Zosteraceae, Cymodoceaceae, Naiadaceae, and Liliaceae by N. TAYLOR; Scheuchzeriaceae by N. L. BRITTON; Alismaceae by J. K. SMALL; Butomaceae, Poaceae (pars) by G. V. NASH. New species are described in the following genera: Sparganium (2), Echinodorus (1), and Trachypogon (2). One new genus (*Machaerocarpus*) of the Alismaceae is proposed, being based on *Damasonium californicum* Torr.—J. M. GREENMAN.

Diseases of trees.—A bulletin, embodying the results of a number of years' investigation of some of the more important diseases of deciduous trees by VON SCHRENK and SPAULDING, has been issued by the national Bureau of Plant Industry.¹⁰ The bulletin contains a large amount of information (there is unfortunately little available) on diseases due to environment, to wound fungi (far

⁷ MOLISCH, H., *Das Warmbad als Mittel zum Treiben der Pflanzen*. 8vo. pp. 38. figs. 12. Jena: Gustav Fischer. 1909. *M* 1.20.

⁸ NAKAI, T., *Flora Koreana. Pars Prima*. Jour. Sci. Coll. 26: 1–304. pls. 1–15. 1909.

⁹ *North American Flora*, Vol. XVII, Part I, pp. 1–98. New York Botanical Garden, 1909.

¹⁰ SCHRENK, H. VON, AND SPAULDING, P., *Diseases of deciduous forest trees*. Bur. Pl. Ind. U. S. Dept. Agric. Bull. 149. pp. 85. pls. 10. figs. 11. 1909.